

Taq DNA Polymerase (5 U/μL)

Product Description

Taq DNA Polymerase is a thermostable DNA Polymerase that catalyzes 5'→3' DNA synthesis. Taq DNA Polymerase has 5'→3' exonuclease activity and is deficient in 3'→5' exonuclease activity. Taq DNA polymerase amplifies uracil-containing templates, can incorporate modified bases, and performs A-tailing on DNA products. This enzyme is provided in a non-hot start format.

Product Applications*

- PCR amplification of DNA fragments (≤5 kb)
- Probe and intercalating dye-based qPCR
- RT-qPCR
- Pre-ligation (dA tailing)
- Homebrew library prep

*Watchmaker Genomics has not tested or validated Taq DNA Polymerase in all applications listed.

Unit Definition and Buffer Composition

- One unit of Taq DNA polymerase is the amount of enzyme that will incorporate 10 nmol of dNTP into activated calf thymus DNA in 30 minutes at 75°C
- Enzyme Storage Buffer: 20 mM Tris-HCl, pH 7.4, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.5% Tween 20
- Standard 10X reaction buffer (not supplied with Kit): 200 mM Tris-HCl, pH 8.3, 400 mM KCl

Kit Contents

Kit	Kit Code	Description	Component Volumes	
			500 μL (2.5 kU)	1000 μL (5 kU)
Taq DNA Polymerase (5 U/μL)	7K0074-500UL 7K0074-1ML	Taq DNA Polymerase (5 U/μL)	500 μL	1000 μL

For larger volumes, higher concentrations, and custom formats, contact the **Sales Team** at sales@watchmakergenomics.com.

Table of Contents

Product Description	1
Product Applications	1
Unit Definition and Buffer Composition	1
Storage and Handling	1
Kit Contents	1
Recommended Reaction Setup	2
Recommended Cycling Protocol	2
Revision History	2

Storage and Handling

Taq DNA Polymerase Kits are shipped on ice packs. Upon receipt, store all kit components at -25°C to -15°C. Keep all components and reaction mixes on ice or a cooled reagent block during routine use. Take care to homogenize solutions thoroughly before use and during reaction setup. Do not vortex the polymerase. When stored and handled as indicated, the product will retain full performance until the expiry date printed on the kit box.

Recommended Reaction Setup

Component	Final Concentration	Volume (per 25 μL reaction)
Buffer (10X) ^{1,2}	1X	2.5 μL
MgCl ₂ (25 mM) ^{1,3}	2 mM	Variable
Each primer 10 μM (Fwd/Rev)	0.2 μM	0.5 μL
Enzyme (5 U/μL) ⁴	0.03 – 0.5 U/μL	Variable
dNTPs (10 mM each)	0.2 mM	0.5 μL
Template DNA ⁵	Genomic 1 – 50 ng	Variable
	Plasmid and viral 1 pg – 1 ng	
PCR grade water	–	Up to 25 μL

¹Buffer (10X) and MgCl₂ are not supplied with Watchmaker Taq DNA Polymerase Kits.

²Watchmaker Taq DNA Polymerase performs well in a range of buffers. The recommended 10X buffer composition is a standard PCR buffer and will need to be optimized for specific applications.

³2 mM is the suggested final MgCl₂ concentration. Mg²⁺ concentrations can alter reaction performance and may need to be optimized. A scouting range between 1 – 6 mM MgCl₂ is recommended. Multiplex and probe-based applications will benefit from increased MgCl₂ concentration.

⁴Enzyme concentration is an important parameter affecting performance and the optimal concentration will vary depending on the application. Optimizing the enzyme concentration is recommended. A good general purpose starting point is 0.05 U/μL. Use higher concentrations of polymerase, up to 0.5 U/μL, in reactions with high concentrations of inhibitors, for fast PCR, or when no amplification is observed. Use lower concentrations of polymerase if non-specific amplification is observed.

⁵Template DNA input is assay dependent and will have to be optimized.

Note: reverse transcriptases are usually inhibitory to PCR enzymes and 1-step qRT-PCR assays may benefit from adding higher concentrations of polymerase.

Recommended Cycling Protocol¹

Step	Temperature (°C)	Time (sec)	Cycles
Initial denaturation	95	60	1
Denaturation	95	15	25 – 35 ⁴
Annealing ²	T _m ~ (T _m - 5)	15	
Extension ³	72	30	
Final extension	72	60	1
–	12	Hold	–

¹Increased PCR specificity can be achieved by performing the reaction setup on ice and inserting reaction tubes into a thermocycler preheated to the denaturing temperature of 95°C

²The optimal annealing temperature should be determined empirically in an annealing temperature gradient experiment.

³The general recommendation for Taq DNA polymerase is 1 min/kb with a minimum of 30 sec for amplicons shorter than 500 bp. Reactions requiring fast PCR can be optimized by increasing the enzyme concentration, shortening the length of the extension step and by performing two-step PCR (combining annealing/extension at the temperature of annealing).

⁴Cycle number will vary depending on starting DNA input.

Revision History

Version	Description	Date
1.0	• First protocol release	12/2023



5744 Central Avenue, Suite 100
Boulder, CO 80301
www.watchmakergenomics.com

For Research Use Only. Not for use in diagnostic procedures.

This content is covered by patents, trademarks, and/or copyrights owned or controlled by Watchmaker Genomics, Inc. For more information, please visit watchmakergenomics.com/licenses. The use of these products may require you to obtain additional third party intellectual property rights for certain applications.

For Technical Support, please contact
support@watchmakergenomics.com.